Geophysical Research Abstracts, Vol. 10, EGU2008-A-04431, 2008 SRef-ID: 1607-7962/gra/EGU2008-A-04431 EGU General Assembly 2008 © Author(s) 2008



## **MESSENGER** observations of Mercury's bow shock and magnetopause

**J. A. Slavin** (1), M. H. Acuña (2), B. J. Anderson (3), D. N. Baker (4), M. Benna (2), S. A. Boardsen (1), G. Gloeckler (5), R. E. Gold (3), G. C. Ho (3), H. Korth (3), S. M. Krimigis (3), S. A. Livi (6), R. L. McNutt Jr. (3), J. M. Raines (5), M. Sarantos (1), D. Schriver (7), S. C. Solomon (8), P. Travnicek (9), and T. H. Zurbuchen (5) (1) Heliophysics Science Division, NASA GSFC, Greenbelt, MD 20771, USA, (2) Solar System Exploration Division, NASA GSFC, Greenbelt, MD 20771, USA, (3) The Johns Hopkins University Applied Physics Laboratory, Laurel, MD 20723, USA, (4) Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA, (5) Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann Arbor, MI 48109, USA (6) Southwest Research Institute, San Antonio, TX 28510, USA, (7) Institute for Geophysics and Planetary Physics, University of California, Los Angeles, CA 90024, USA, (8) Department of Terrestrial Magnetism, Carnegie Institution of Washington, DC 20015, USA, and (9) Institute of Atmospheric Physics, Prague, Czech Republic, 14131

MESSENGER's 14 January 2008 encounter with Mercury will provide the first new observations of the solar wind interaction with this planet since the Mariner 10 flybys that took place over 30 years ago. The closest approach distance for this first MES-SENGER flyby is targeted for an altitude of 200 km as compared with the 707 km and 327 km attained by Mariner 10 on 29 March 1974 and 16 March 1975, respectively. The locations of the bow shock and magnetopause boundaries observed by MESSEN-GER will be examined and compared against those found in the earlier Mariner 10 measurements and the predictions of theoretical models and numerical simulations. The structure of the magnetopause will be investigated for the presence of flux transfer events or other evidence of magnetic reconnection as will the more general implications of these new MESSENGER bow shock and magnetopause observations for the global solar wind interaction with Mercury.