

# Hidden Markov models for linking synoptic atmospheric patterns to daily precipitation in Bulgaria

N. Neykov (1), W. Zucchini (2), P. Neytchev (1) and H. Hristov (1)

(1) National Inst. of Meteorology and Hydrology, Bulgarian Academy of Sciences, Sofia, Bulgaria, (2) Institute of Statistics and Econometrics, University of Göttingen, Göttingen, Germany (neyko.neykov@meteo.bg)

The non-homogenous hidden Markov Models (NHMM) have found widespread application in meteorology and hydrology in Australia, North and South America, e.g., in studies of climate variability or climate change, statistical downscaling of daily precipitation from observed and numerical climate model simulations. Details can be found in Zucchini and Guttorp (1991), Hughes and Guttorp (1994), Hughes et al. (1999), Charles et al. (1999), Bellone et al. (2000), Charles et al. (2003) and Charles et al. (2004), Robertson et al. (2004), just to name a few. However, the NHMM methodology still has not been employed for similar purposes in Europe. The NHMM links large-scale atmospheric patterns to daily precipitation data at a network of rain gauge stations, via several hidden (unobserved) states called the “weather states”. The evolution of these states is modeled as a first-order Markov process with state-to-state transition probabilities conditioned on some indices of the atmospheric variables. Due to these weather states the spatial precipitation dependence can be partially or completely captured, see Hughes et al. (1999). In the present study various NHMMs are used to relate daily precipitation at 30 rain gauge stations covering broadly the territory of Bulgaria to synoptic atmospheric data. A 40-year record (1960-2000) of daily winters (October through March) precipitation amounts is used at each station. The atmospheric data consists of daily sea-level pressure, geopotential height at 500 hPa, air temperature at 850 hPa and relative humidity at 700 hPa on a  $2.5^\circ \times 2.5^\circ$  grid based on NCEP-NCAR reanalysis dataset covering the Europe-Atlantic sector  $30^\circ\text{W}$ – $60^\circ\text{E}$ ,  $20^\circ\text{N}$ – $70^\circ\text{N}$  for the same period. The first 30 years data are used for model fitting purposes while the remaining 10 years are used for model evaluation. Detailed model validation is carried out on various aspects. The identified weather states are found to be physically interpretable in terms of regional climatology.

## References

Bellone,E., Hughes,J.P. and Guttorp,P. (2000) A hidden Markov model for relating synoptic scale patterns to precipitation amounts. *Climate Research*, 15, 1-12.

Hughes,J.P. and Guttorp,P. (1994). A class of stochastic models for relating synoptic atmospheric patterns to regional hydrologic phenomena. *Water Resour. Res.* 30, 1535-1546.

Hughes,J.P., Guttorp P. and Charles,S. (1999). A nonhomogeneous hidden Markov model for precipitation occurrence. *J R. Stat. Soc. Ser C*, 48, 15-30.

Charles,S., Bates,B., and Hughes,P. (1999). A spatio-temporal model for downscaling precipitation occurrence and amounts. *J. Geoph. Res.*, 104, 31657-31669.

Charles,S., Bates,B. and Vilney,N. (2003). Linking atmospheric circulation to daily rainfall patterns across the Murrumbidgee River Basin, *Water Science and Technology*, 48, 223-240.

Charles,S. Bates,B., Smith,I. and Hughes,P. (2004). Statistical downscaling of daily precipitation from observed and modeled atmospheric fields. *Hydrological Process*, 18, 1373-1394.

Zucchini,W. and Guttorp,P. (1991). A hidden Markov model for space-time precipitation. *Water Resour. Res.*, 27,1917-1923.

MacDonald,I. and Zucchini,W. (1997). *Hidden Markov and other models for discrete-valued time series models*. Chapman & Hall, London.

Robertson,A.W., Kirshner,S. and Smyth,P. (2004). Downscaling of daily rainfall occurrence over Northeast Brazil using a Hidden Markov Model, *J. Climate*, 17, 4407-4424.