Geophysical Research Abstracts, Vol. 7, 06144, 2005 SRef-ID: 1607-7962/gra/EGU05-A-06144 © European Geosciences Union 2005



Satellite laser ranging and the terrestrial reference frame

J. Ries (1), M. Cheng (1), D. Chambers (1)

(1) Center for Space Research, The University of Texas at Austin (ries@csr.utexas.edu)

The importance of an accurate and stable reference frame, for reliable interpretation of signals from the various Earth system research areas such as post-glacial rebound and sea level change, cannot be overstated. The various geodetic observing systems are at the core of the determination and maintenance of the celestial and terrestrial reference frames on which other sciences depend, and each system provides essential and unique contributions. Satellite laser ranging (SLR) to geodetic satellites has provided long-term observations of the long-wavelength temporal mass redistribution in the Earth system and is especially important for the determination of the absolute geocenter location, an essential component of the reference frame definition. Most uniquely, the determination of the most fundamental gravity parameter, the gravitational coefficient of the Earth (GM), is still best determined through the analysis of SLR tracking data. The accurate determination of GM is critical in the definition of the absolute scale of the geocentric reference frame This paper examines the state of our knowledge of the above components, especially GM and geocenter, as well as an assessment of the sensitivity to recent changes in the SLR network.