



Segregation in binary mixtures

M. Larcher (1), B. Spinewine (2), L. Fraccarollo (1) & A. Armanini (1)

(1) CUDAM - Dipartimento di Ingegneria Civile e Ambientale - Università di Trento, Italy, (2)
Department of Civil and Environmental Engineering - Université catholique de Louvain,
Belgium

To investigate the vertical structure of free-surface liquid-granular flows, it is of particular interest to be able to materialise steady flow conditions. With this experimental technique, in the recent past, a thorough activity has been devoted to the rheology of flowing mixtures of well-sorted granular particles and water (Armanini et al., 2005). In this contribution, we present a preliminary analysis of the evolution, in time and space, of bimodal mixtures in quasi-uniform conditions. In accordance with experimental evidence given by Savage & Lun (1988), Gray & Hutter (1997) and Gray (2001), segregation of particles of different sizes has been observed, with the formation of layers with the largest particles on top and the smaller at the bottom. This phenomenon has been investigated with the support of a newly developed imaging technique, based on the development of findings by Spinewine et al. (2003), able to tackle non-uniformly sized granular materials (within IMPACT European Project, 2001-2004). Moreover, a special device was built to allow raw measurements of the distribution of the relative solid-concentration of the two size fractions. Therefore, data from sidewall image and bulk measurements allow also to infer some information about three dimensional effects. The larger sediments employed are PVC particles with an equivalent-spherical diameter equal to 3.7 mm, also utilized for the experiments with a single size-fraction, while the smaller are quite uniform plastic spheres having a mean diameter of 0.9 mm. Both materials have a density of about 1570 kg/m³. In the experiments, a mixture of water and sediments was introduced steadily in the upstream part of the flume in well-mixed conditions, with a quite uniform distribution of both granulometric classes throughout the flow depth. A progressive particle segregation was observed in downstream direction, with the larger particles migrating upwards and the smaller ones downwards. The distance measured experimentally at which segregation is almost complete was compared with the values predicted by the theory of Vallance &

Savage (2000). Theoretical results can be considered results in good accordance with experimental evidence.

REFERENCES ARMANINI, A., CAPART, H., FRACCAROLLO, L., LARCHER, M. (2005). Rheological stratification in experimental free-surface flows of granular-liquid mixtures. *J. Fluid Mech.*, in press. GRAY J.M.N.T. & HUTTER K. (1997) Pattern formation in granular avalanches. *Continuum Mechanics and Thermodynamics* 9, 341-345. GRAY J.M.N.T. (2001) Granular flow in partially filled slowly rotating drums. *J. Fluid. Mech.* 441, 1-29. SAVAGE S.B. & LUN C.K.K. (1988) Particle size segregation in inclined chute flow of dry cohesionless granular solids. *Journal of Fluid Mechanics* 189, 311-335. SAVAGE S.B. & HUTTER K. (1989) The motion of a finite mass of granular material down a rough incline. *Journal of Fluid Mechanics* 199, 177-215. SPINEWINE B., CAPART H., LARCHER M., ZECH Y. (2003). Three-dimensional Voronoï imaging methods for the measurement of near-wall particulate flows. *Experiments in Fluids*, vol. 34, pp. 227-241. VALLANCE J.W. & SAVAGE S.B. (2000) Particle segregation in granular flows down chutes, *IUTAM Symposium on segregation in granular flows*, 31-51.