



NorthGRIP – a Greenland ice core reaching back into the Eemian Period

D. Dahl-Jensen (1), J. P. Steffensen (1), S. J. Johnsen (1), J. Jouzel (2), J. White (3), H. Miller (4), J. Schwander (5), M. Hansson (6), Th. Thorsteinsson (7), Y. Fujii (8), R. Lorrain (9)

(1)Niels Bohr Institute for Astronomy, Physics and Geophysics, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen OE, Denmark, (2)Institute Pierre Simon Laplace/ Laboratoire des Sciences du Climat et de l'Environnement, UMR CEA-CNRS, CE Saclay, Orme des Merisiers, 91191 Gif-Sur-Yvette, France, (3) INSTAAR, Campus Box 450, University of Colorado, Boulder, CO 80309-0450, USA, (4) Alfred-Wegener-Institute for Polar- and Marine Research (AWI), Postfach 120161, D-27515 Bremerhaven, Germany, (5) Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, CH-3012, Switzerland, (6) Department of Physical Geography and Quaternary Geology, Stockholm University, S-106 91, Stockholm, Sweden, (7) National Energy Authority, Grensásvegur 9, IS-108 Reykjavík, Iceland, (8) National Institute of Polar Research, Kaga 1-9-10, Itabashi-ku, Tokyo 173-8515 Japan, (9) Département des Sciences de la terre et de l'Environnement, Faculté des Sciences, CP 160/03, Université Libre de Bruxelles, 50 avenue FD Roosevelt, B1050 Brussels, Belgium, (ddj@gfy.ku.dk / Fax: +45 35 36 53 57 / Phone: +45 35 32 05 56)

While the Greenland ice cores have a very high resolution the ice cores available at present only reach 125.000 years back in time thus reaching into the last interglacial period, the Eemian. It is a very high wish in the ice core and climate community to obtain an ice core from Greenland reaching further back in time.

Based on the net of Radio Echo Sounding (RES) profiles over the Greenland ice sheet and the calibration of the age of the observed internal layers at the deep drill sites GRIP and NGRIP an inverse model has been developed to date the ice at locations where internal RES layers are detectable.

A location at (75.5N, 46W) has been selected as a good site. The ice thickness is 2550 m, present accumulation rate is 0.23 m ice equiv./yr and pre-Eemian ice is expected to

be found in the basal 200 m of ice.

Besides from being a site with old basal ice the site is also selected as a promising site for studies of the present interglacial climate covering the last 10.000 years because the precipitation is more single sourced than that found at the other deep drilling sites in Greenland. The isotopic signals are thus expected to be more directly related to temperatures changes during this period.