



## **Electromagnetic Ion Cyclotron Waves, Plasma Plumes and Cold Ion Density Measurements in the Magnetosphere**

**B. J. Fraser** (1), H. J. Singer (2), J. Goldstein (3), D. L. Gallagher (4), and M. Thomsen (5)

(1) CRC for Satellite Systems, School of Mathematical & Physical Sciences, University of Newcastle, NSW 2308, Australia, (brian.fraser@newcastle.edu.au / Phone +61 2 4921 5445)

(2) NOAA/SEC, 325 Broadway, Boulder, CO 80305, USA

(3) Department of Physics and Astronomy, Rice University, Houston, Texas, USA

(4) Space Science Department, Marshall Space Flight Center, National Space Science & Technology Center, Huntsville, AL 35805

(5) Los Alamos National Laboratory, Los Alamos, New Mexico, USA

The radial plasma density profile in the plasmasphere-plasmapause region may vary considerably depending on the particle species measured and the particular satellite instrument making the measurement. For example the heavy ion relative concentrations of  $\text{He}^+$  and  $\text{O}^+$  compared with  $\text{H}^+$  ions, are not constant with increasing radial distance. Also, some satellites measure electron density, directly or indirectly, while others measure  $\text{H}^+$  and  $\text{He}^+$ . IMAGE-EUV satellite results have confirmed the existence of plasma plumes in the middle magnetosphere and often electromagnetic ion cyclotron waves are observed within these plumes. This paper will introduce the relationship between plumes and EMIC waves. It will then consider intercalibration between instruments measuring plasma density in the plasmasphere/magnetosphere and develop diagnostic techniques to compute heavy ion densities in plasma plumes. Using the spectral properties of EMIC waves seen in plumes in conjunction with ion density data and diagnostic techniques it is possible to almost completely describe the plasmasphere and magnetosphere cold/cool plasma composition. Particle data from the LANL geostationary satellites, EUV images from the IMAGE satellite and EMIC

wave data from the GOES satellites are used in this study. Results yield ion densities ranging from over  $H^+ = 30 - 80 \text{ cm}^{-3}$  and  $He^+ = 7 - 13 \text{ cm}^{-3}$  and  $O^+ = 5 - 9 \text{ cm}^{-3}$ . The  $He^+$  results are compared with computed pseudo-densities of  $He^+$  determined from IMAGE-EUV data.