



## **A unique carbon isotope record across the Guadalupian-Lopingian (Permian) boundary in Panthalassan paleoatoll carbonates: the high-productivity “Kamura event” and its collapse**

**Y. Isozaki** (1), H. Kawahata (2) and A. Ota (1)

1. Dept. Earth Sci. & Astron., Univ. Tokyo, Japan (isozaki@chianti.c.u-tokyo.ac.jp)
2. Ocean Research Inst., Univ. Tokyo, Japan

Middle to Upper Permian shallow marine carbonates in Kamura, SW Japan, were derived from a paleo-atoll complex on a paleo-seamount in mid-Panthalassa. A secular change in  $\delta^{13}\text{C}_{carb}$  was analyzed to document the oceanographic change in the superocean Panthalassa with respect to the mass extinction across the Guadalupian-Lopingian boundary (G-LB). The Upper Guadalupian is characterized mostly by unusually high positive values of +4.9 to +6.2 per mil, whereas the Lower Lopingian by low values from +1.9 to +3.5. The 4 per mil negative shift in total occurred in 3 steps around the G-LB. A remarkably sharp 2.4 per mil drop occurred in the topmost 2 m interval of the Guadalupian, after large fusulines disappeared abruptly. Such a prominent high positive plateau followed by a large negative shift across the G-LB was first detected in the mid-superocean and is correlated with the stratotype of the G-LB in China. The present results prove that the end-Guadalupian event occurred doubtlessly on a global scale. The end-Guadalupian high positive plateau in  $\delta^{13}\text{C}_{carb}$  values over +5 per mil is particularly noteworthy because it recorded an unusually high bio-productivity period that has not been described in the Permian. This end-Guadalupian high productivity event, newly named Kamura event, suggests burial of a huge amount of organic carbon, draw-down of atmospheric CO<sub>2</sub> and resultant global cooling in the late Guadalupian, considerably after the Gondwana glaciation. The low temperatures during the Kamura event may have caused the end-Guadalupian extinction of large-shelled Tethyan fusulines adapted to warm climate. The Kamura event agrees with the global sea-level curve in the Middle-Late Permian.