



MAP3D: A MESOSCALE AIR POLLUTION 3 D MODELLING TOOL

O. Couach (1, 2), F. Kirchner (1), I. Balin (1), P. Ristori (3), V. Simeonov (3), and H. van den Bergh (3)

(1) EnviroScopY SA (start-up project), PSE - EPFL, CH-1015 Lausanne, Switzerland,

(2) Swiss Federal Institute of Technology, ENAC/ISTE/EFLUM, CH 1015 Lausanne, Switzerland

(3) Swiss Federal Institute of Technology, ENAC/ISTE/LPAS, CH 1015 Lausanne, Switzerland

Contact: olivier.couach@epfl.ch , Phone: +41 21 693 6139

The assessment of the regional (e.g. mesoscale) air pollution problematic needs the development of physical-chemical based model instead of dispersion (transport) based models.

A relatively new mesoscale air pollution modeling approach, integrating new modules for impact studies and forecast) namely **MAP3D** is presented here. The potential of realistic, calibrated, validated and thus representative outputs will be illustrated for different pollutants as O_3 , NO_x , CH_2O , CO and meteorological parameters in selected field campaigns (e.g. Grenoble, Strasbourg and Swiss Plateau).

Comparisons and validation with representative 3D measurements (DOAS, LIDAR, radio- soundings, wind-profilers and aircraft) and results obtained by a combined 3D modeling - measurements approach will be shown. The item of air pollution regimes (NO_X/VOC) and specific indicators will be also addressed.

Finally the capability of MAP3D for realizing representative studies, impact and scenarios and air pollution forecast as a policy maker decision tool will be discussed.

Keywords: physical-chemical based modeling, 3D, mesoscale, air pollution, photo-chemistry, ozone, indicators, VOC, NOX, planetary boundary layers dynamics