



## What is the strength of ‘CO<sub>2</sub>-calcification’ feedback on future fossil fuel CO<sub>2</sub> uptake?

**A. Ridgwell** (1), I. Zondervan (2), J. Hargreaves (3), T. Lenton (4), and J. Bijma (5)

(1) Department of Earth and Ocean Sciences, University of British Columbia, Vancouver BC, Canada [andy@seao2.org], (2) Biogeosciences, Alfred Wegener Institute for Polar- and Marine Research, Am Handelshafen 12, Bremerhaven, Germany [izondervan@awi-bremerhaven.de], (3) Frontier Research Center for Global Change, 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Kanagawa 236-0001, Japan [jules@jamstec.go.jp], (4) School of Environmental Sciences, University of East Anglia, Norwich, UK [T.Lenton@uea.ac.uk], (5) Biogeosciences, Alfred Wegener Institute for Polar- and Marine Research, Am Handelshafen 12, Bremerhaven, Germany [jbijma@awi-bremerhaven.de].

Understanding the strength of the ‘CO<sub>2</sub>-calcification’ feedback; the enhancement of fossil fuel CO<sub>2</sub> uptake by the ocean due to a reduction in marine carbonate production, is critical if we are to accurately predict the future trajectory of atmospheric CO<sub>2</sub>. To do this, computer models of the ocean carbon cycle have typically been informed by the *in vitro* response of the coccolithophorid *Emiliana huxleyi* to a decrease in ambient carbonate ion concentration. However, analysis of the experiments carried out to date across a range of calcifying plankton species leads us to suspect that the CO<sub>2</sub>-calcification feedback could be much more important than previously assumed. Furthermore, analysis of the anthropogenic CO<sub>2</sub> impact on marine calcification has yet to include the effect of synchronous changes in climate. We address these issues and the uncertainties surrounding the strength of the CO<sub>2</sub>-calcification feedback with the aid of an ensemble of instances of the GENIE-1 Earth system model. We find that an additional ca. 200 PgC could be sequestered in the ocean by the end of this millennium.