



## Estimating lightning produced $\text{NO}_x$ from satellite

S. Beirle (1), U. Platt (1), T. Wagner (1)

(1) IUP Uni Heidelberg, INF 229 69120 Heidelberg

Nitrogen oxides ( $\text{NO}_x = \text{NO} + \text{NO}_2$ ) play an important role in tropospheric chemistry, in particular in catalytic ozone production. Lightning provides a natural source of nitrogen oxides. Recent estimates of lightning produced  $\text{NO}_x$  ( $\text{LNO}_x$ ) are of the order of 5 Tg [N] per year with still high uncertainties in the range of one order of magnitude.

Satellite sensors like GOME or SCIAMACHY allow the retrieval of tropospheric vertical column densities (TVCDs) of  $\text{NO}_2$  on a global scale. Recently, correlations of enhanced  $\text{NO}_2$  TVCDs with lightning have been reported.

Here we discuss the potential of satellite measurements of  $\text{NO}_2$  for the quantitative estimation of lightning produced  $\text{NO}_x$ . The sensitivity (AMF) of satellite instruments to lightning  $\text{NO}_x$  and the impact of additionally required information (e.g. flash rates,  $\text{NO}_x$  profile,  $\text{NO}_x$  partitioning,  $\text{NO}_x$  lifetime, transport) are analyzed in detail.

We present the results of quantitative  $\text{LNO}_x$  estimates for statistical approaches as well as for some individual case studies, in particular a lightning event in the Gulf of Mexico, coinciding with the GOME measurement in space and time.