



One-to-one interhemispheric coupling of multicentennial glacial climate variability recorded in the EPICA DML ice core

H. Fischer and the EPICA community

Alfred-Wegener-Institute for Polar and Marine Research, Columbusstrasse, 27568 Bremerhaven, Germany, (hufischer@awi-bremerhaven.de)

Precise knowledge of the phase relationship between climate changes in both hemispheres is a key to understand the coupling mechanisms in the Earth climate system. The glacial and deglacial climate evolution in the North Atlantic region was characterized by rapid shifts from cold stadial to warmer interstadial conditions [NorthGRIP community members, 2004]. In contrast climate in the Circum-Antarctic region as recorded in ice cores from the Indian and Pacific Ocean sector of the Antarctic ice sheet [Blunier and Brook, 2001; EPICA community members, 2004; Watanabe et al., 2003] exhibited slower millennial changes with temperature amplitudes of up to 3 °C during that time. Absolute synchronization of Greenland and Antarctic ice core records [Blunier and Brook, 2001] using the global atmospheric change in CH₄ concentrations accompanying the DO events in the North showed a conspicuous phase relationship with the South starting to cool as soon as the longest DO events in the North set in. This bipolar seesaw pattern can be explained by the transport of heat and freshwater connected to the Atlantic Meridional Overturning Circulation (AMOC) [Knutti et al., 2004; Stocker and Johnsen, 2003], where a stronger (reduced) AMOC leads to increased (decreased) drainage of heat from the Southern Ocean heat reservoir. To what extent this concept is also able to explain the shorter and less pronounced climate variability found in Antarctic ice cores, however, remained obscure. Here we report on the high-resolution climate record from the new EPICA Dronning Maud Land (EDML) ice core which represents the first direct South Atlantic counterpart to the Greenland records. Methane synchronized, high resolution EDML isotope temperatures ($\delta^{18}\text{O}$) clearly show a one-to-one bipolar seesaw coupling also for all the shorter DO events, with the amplitude of the corresponding Antarctic warming being

dependent on the duration of the concurrent stadial in the North.

Blunier, T., and E. J. Brook (2001), Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, *Science*, 291, 109-112.

EPICA community members (2004), Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623-628.

Knutti, R., J. Flückiger, T. F. Stocker, and A. Timmermann (2004), Strong hemispheric coupling of glacial climate through freshwater discharge and ocean circulation, *Nature*, 430, 851-856.

North Greenland Ice Core Project members (2004), High resolution climate record of the northern hemisphere reaching into the last interglacial period. *Nature* 431, 147-151.

Stocker, T., and S. Johnsen (2003), A minimum thermodynamic model of the bipolar seesaw, *Paleoceanogr*, 18, (4), doi:10.1029/2003PA000920.

Watanabe, O., J. Jouzel, S. Johnsen, F. Parrenin, H. Shoji, and N. Yoshida (2003), Homogeneous climate variability across East Antarctica over the past three glacial cycles, *Nature*, 422, 509-512.