



## **ACE (Atmospheric Chemistry Experiment) Measurements of the upper troposphere and stratosphere**

**C. Rinsland (1)**

(1) NASA Langley Research Center, Mail Stop 401A, Hampton, VA 23681-2199 U.S.A.

We highlight chemistry and trend measurements from the Atmospheric Chemistry Experiment (ACE) which is providing precise middle troposphere to the lower thermosphere measurements of the Earth's upper troposphere to lower mesosphere with the solar occultation technique. The primary instrument is a 0.02 cm<sup>-1</sup> resolution Fourier transform spectrometer covering 750-4400 cm<sup>-1</sup> simultaneously. ACE also known as SCISAT 1 is a Canadian lead mission launched into a 74° inclined orbit at 650 km altitude on August 12, 2003 by a U.S.-supplied Pegasus vehicle. Additional measurements are obtained with a UV-visible spectrometer and two imagers providing measurements at 0.525 and 1.02 microns. The Sun produced some of the most powerful solar flares ever recorded between 18 October and 5 November 2003. ACE measurements of Arctic NO<sub>x</sub> vs. pressure from 0.07-2 hPa (~40-60 km) during February and March 2004 showed the dilution of NO<sub>x</sub> resulting from photochemical loss in the sunlit atmosphere, and the time series of measurements were used to estimate the descent rate from the upper stratospheric NO<sub>x</sub> time series. The capability of solar occultation instruments to infer nearly all components of the stratospheric inventories of chlorine and fluorine from high spectral resolution solar occultation instruments was pioneered by the Atmospheric Trace Molecule Spectroscopy (ATMOS) Fourier transform spectrometer. The ACE instrument improved on the ATMOS design, and has been used to infer the trends of both chlorine and fluorine at a critical time when stratospheric chlorine is near its maximum. Trend studies comparing ATMOS and ACE measurements have been reported in addition to the detections of HCFC-142b and CFC-113. Inventories for chlorine and fluorine in 2004 have been inferred including the detection of CO-CIF, a missing component in the chlorine and fluorine chemical families from coadded ACE spectra in the region of its strongest infrared band. Frequent clear sky scenes oc-

cur in the middle and upper troposphere. ACE measured several molecular species not previously detected from space. Important new results are the detection of methanol (CH<sub>3</sub>OH), the most abundant organic molecule in the troposphere, and formic acid (HCOOH). ACE measures southern hemisphere lower to mid-latitudes during August to October when intense vegetative fires occur in tropical regions of Africa and South America. Convection from those fires results in transport of the pollutant emissions to the middle and upper troposphere, where molecules with a sufficiently long lifetime can be measured by ACE. Measurements include formic acid (HCOOH) an important oxygenated volatile organic compound (OVOC) with major limitations recognized in the ability of models to reproduce OVOC chemistry, particularly in the dry upper troposphere where OVOCs are a major source of HO<sub>x</sub> in the background troposphere with ACE measurements showing elevated HCOOH in upper tropospheric plumes during both 2004 and 2005. Additional measurements from plumes include H<sub>2</sub>O<sub>2</sub> (hydrogen peroxide). ACE also measures boreal regions which are frequently impacted during summer by biomass burning. Summer 2004 measurements show elevated upper tropospheric CO, C<sub>2</sub>H<sub>6</sub>, HCN, CH<sub>3</sub>Cl, CH<sub>4</sub>, C<sub>2</sub>H<sub>2</sub>, CH<sub>3</sub>OH, HCOOH, and OCS mixing ratios in plumes up to 189 ppbv (10<sup>-9</sup> per unit volume) for CO, 1.40 ppbv for C<sub>2</sub>H<sub>6</sub>, 830 pptv (10<sup>-12</sup> per unit volume) for HCN, 1.78 ppbv for CH<sub>3</sub>Cl, 2.16 ppmv (10<sup>-6</sup> per unit volume) for CH<sub>4</sub>, 0.118 ppbv for C<sub>2</sub>H<sub>2</sub>, 3.89 ppbv for CH<sub>3</sub>OH, 0.843 ppbv for HCOOH, and 0.543 ppbv for OCS in western Canada and Alaska at 50°N-68°N latitude between 29 June and 23 July 2004. Emission factors have been inferred and compared with previously reported values.