



## **Planetary magnetism driven by convection**

**S. Starchenko**

IZMIRAN, Troitsk, Moscow reg., 142190 Russia (sstarchenko@mail.ru)

Planetary convection and magnetic instabilities driven by thermal or/and compositional power are investigated in their natural limit of very small transport coefficients. Correspondent laboratory convection onset modeling is suggested and theoretically explored up to the asymptotic values suitable for the planets and large moons. For the Earth's type planets/moons the strong influence of the inner rigid core size is found on such double diffusion convection/magnetism onset and support. The relative size about half of the modern Earth's inner core size makes compositional convection valuable and able on addition magnetic support as in the past Earth or on magnetic appearance possible in future Venus. Then the core growing supports thermal convection (and related magnetism, e.g. in Ganymede, if any) up to the about half of the convective shell size. Further growing of the inner core suppresses the compositional convection which died out when thickness of the shell become too small. The last possibly was in the past Mars when its magnetic dynamo has been stopped. Thermal convection alone seems enough to support magnetic fields in the outer planets. The principal MAC balance between Magnetic, Archimedean buoyancy and Coriolis force is about the same in the Earth's, Jupiter and Saturn. Uranus, Neptune and perhaps Ganymede unusual magneto-convection should be in another not yet explored IAC balance that is between Inertia, Archimedean and Coriolis force those exceed magnetic force.