



Evaluation of global cloud regimes in contemporary GCMs

K. D. Williams (1), G. Tselioudis (2)

(1) Hadley Centre, Met Office (keith.williams@metoffice.gov.uk), (2) NASA GISS

The radiative feedback from clouds remains the largest source of variation in climate sensitivity amongst general circulation models (GCMs). A cloud clustering methodology is applied to six contemporary GCMs in order to provide a detailed intercomparison and evaluation of the simulated cloud regimes. By analysing GCMs in the context of cloud regimes, processes related to particular cloud types are more likely to be evaluated. In this study, the mean properties of the global cloud regimes are evaluated, and the cloud response to climate change is analysed in the cloud-regime framework. Most of the GCMs are able to simulate the principal cloud regimes, however none of the models analysed have a good representation of trade cumulus in the tropics. The models also share a difficulty in simulating those regimes with cloud tops at mid-levels, with only ECHAM5 producing a regime of tropical cumulus congestus. Optically thick, high top cloud in the extra-tropics, typically associated with the passage of frontal systems, is simulated considerably too frequently in the ECHAM5 model. This appears to be a result of the cloud type persisting in the model after the meteorological conditions associated with frontal systems have ceased. The simulation of stratocumulus in the MIROC GCMs is too extensive, resulting in the tropics being too reflective.