



Long cores through complete cold-water coral carbonate mounds: from IODP307 to EuroMARC project CARBONATE

A.J. Wheeler (1), A. Freiwald (2), D. Hebbeln (3), R. Swennen (4), T.C.E. Van Weering (5), H. De Haas (5), B. Dorschel (1)

(1) Dept. of Geology & Environmental Research Institute, University College Cork, Cork, Ireland, (2) Institute of Palaeontology, Erlangen University, Erlangen, Germany, (3) MARUM-Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany, (4) Sedimentpetrologie, Leuven Catholic University, Heverlee, Belgium, (5) Royal Netherlands Institute for Sea Research, Texel, The Netherlands, (a.wheeler@ucc.ie / Fax: +353 (0)21 4901932 / Phone: +353 (0)21 4901943)

Along the European Atlantic continental margin, recent to young fossil carbonate mounds and build-ups by cold-water corals (predominantly the framework forming ahermatypic corals *Lophelia pertusa* and *Madrepora oculata*) occur from northern Norway to the Gulf of Cadiz with an emerging global distribution pattern showing a European concentration (see Roberts *et al.*, 2006). Up to now the carbonate stored in carbonate mounds has not been considered in any global carbonate budget or linked to any global carbon budget involving greenhouse gases. A major challenge exists to quantify the amount and flux of carbon stored by these newly discovered areas of enhanced carbonate accumulation in intermediate water depth (e.g. carbonate mounds). The only existing integrative cold-water coral carbonate budget that has been performed is on short, post-glacial cores relevant to the Norwegian shelf only, which suggests that these small cold-water coral carbonate mounds alone may account for an additional 1% of total marine carbonate production (Lindberg *et al.*, 2005).

Furthermore, investigations so far reveal that all mounds possess different growth histories depending on the environmental setting and the involved faunal associations. Unfortunately, existing cores only penetrated the upper few meters of the mounds thus limiting mound research to the very late stage of mound development. Access to the longer sequences preserved in giant carbonate mounds was overcome in May 2005

when the IODP Expedition 307 (Porcupine Mound Drilling) recovered complete sedimentary records from the 155 m high “Challenger Mound” in the Porcupine Seabight west off Ireland, including the underlying strata and the enigmatic mound base. Ship-board and preliminary shore based investigations of the recovered sediments exposed complex internal mound sedimentary structures and sedimentary fabrics. Coral occurrences throughout the core stressed the biological influence on mound development and construction. The common occurrence of hardgrounds is indicative of harsh environmental controls on mound formation and development with diagenetic effects playing a progressive role in internal mound geochemical environment influencing cementation, dissolution, porosity and permeability.

IODP has delivered one complete record from base to top of one carbonate mound in the Belgica Mound Province, Porcupine Seabight, NE Atlantic. EU FP projects have revealed late stage history of giant mounds in different settings showing that different mounds respond in different ways to environmental forcing factors with no one mound being typical of all. By understanding how biogeochemical processes control the development of these carbonate mounds and their response to climate change, we will make an important step in quantifying their role as mid-latitude carbonate sinks. In the end, a better understanding of the processes involved in mound formation and development may also result in new views on fossil analogues many of which are less accessible hydrocarbon reservoirs.