



## The formation of rock glaciers from mass movements

**J.M. Reitner** and A. Gruber

Geological Survey of Austria, Vienna

(juergen.reitner@geologie.ac.at; alfred.gruber@geologie.ac.at)

According to our present knowledge (BARSCH, 1996; HAEBERLI et al. 2006) the morphological and geological conditions for the formation of rock glaciers in Alpine environments seem to be clear. All known examples derive from porous more or less coarse grained sedimentary bodies, either from moraines or, in most cases, from talus fans. In the latter case the debris accumulation originates overwhelmingly from physical weathering, rock falls or rock avalanches in proximity to rockwalls. However, in the course of geological mapping in the crystalline areas of Eastern and Northern Tyrol (Schober Gruppe, Tuxer Alpen) we found an additional setting. Some relict rock glaciers occur directly at the bulging toe of bedrock slopes, which had been affected by deep-seated gravitational slope deformations (REITNER, 2003; GRUBER, 2005). Furthermore rock glaciers are also present in ridge-top depressions and similar graben-like features that originated from gravitational processes in jointed bedrock. In all these cases talus fans with debris accumulation are missing in the source area of those rock glaciers.

According to our model the disintegration of jointed rocks by creeping mass movements resulted in an increased volume of joint space. This enabled the formation of interstitial ice under permafrost conditions. Increased ice saturation led to the reduction of the angle of internal friction and finally to the initial formation of a rock glacier. Due to the previous and maybe still ongoing slope deformation abundant material was provided for the further movement and thus for formation of quite large rock glaciers. Most rock glaciers of this type originated from mass movements of sagging -type (Sacking sensu ZISCHINSKY, 1966), which illustrates the continuous transition from gravitational to periglacial creep process in high Alpine areas.

All studied examples are of Lateglacial age according to the altitude in correspondence

to the known amount of permafrost depression compared to modern time. Thus, on the one hand such rock glaciers postdate the formation of the mass movements, which enable a chronological constraint of this phenomenon on the base of our knowledge of climate history. On the other hand, those examples with rock glaciers linked at various altitudes with mass movements mirror as well former stepwise permafrost degradation, where rock glacier formation moved to higher altitudes. In this respect and envisaging a rising permafrost boundary, rock glacier formation at slopes affected by mass movements should be anticipated for the future.

#### References:

BARSCH, D. (1996): Rockglaciers. - Springer Verlag, Berlin.

GRUBER, A. (2005) Bericht 2004 über geologische Aufnahmen im Quartär der Nördlichen Tuxer Alpen auf Blatt 148 Brenner.- Jahrbuch der Geologischen Bundesanstalt, 145, 337-343, Wien.

HAEBERLI, W. et al. (2006): Permafrost Creep and Rock Glacier Dynamics.- Permafrost and Periglac. Process., 17, 189-214 (2006), Wiley Interscience, New York

REITNER, J. M. (2003a): Bericht 1998-99 über geologische Aufnahmen im Quartär und Kristallin auf Blatt 179 Lienz.- Jahrbuch der Geologischen Bundesanstalt., 143, 514-522, Wien.

ZISCHINSKY, U. (1966): On the deformation of high slopes. Proc.-1st Int. Conf.Soc.Rock Mech. Lisbon, 179-185.