



## Saturn's Titan: Evidence for surface activity

R. M. Nelson(1), L. Kamp(1), D. L. Matson(1), P. G. J. Irwin(2), K. H. Baines(1), M. D. Boryta(3), F. E. Leader (1), R. Jauman(4), W. D. Smythe(1), C. Sotin(5), R. N. Clark(6), D. P. Cruikshank(7), P. Drossart(8), J. C. Pearl (9), B. W. Hapke(10), J. Lunine(11), M. Combes (12), G. Bellucci(13), J.-P. Bibring(14) F. Capaccioni(13), P. Cerroni(13), A. Coradini (13) V. Formisano(13), G. Filacchione(13) R. Y. Langevin(14), T. B. McCord (15), V. Mennella(16), P. D. Nicholson (17), B. Sicardy(8)

(1)JPL/NASA, Pasadena, CA USA, (2)Atmospheric, Oceanic and Planetary Physics, Clarendon Laboratory, Parks Road, Oxford, UK, (3)Mount San Antonio College, Walnut, CA USA, (4)Institute for Planetary Exploration, DLR, Berlin, Germany, (5)University of Nantes, Nantes, France, (6)USGS, Denver, CO, USA, (7)NASA AMES, Mountain View, CA (8)Observatoire de Paris-Meudon, France, (9)Goddard Space Flight Center, Greenbelt MD, (10)U of Pittsburgh, Pittsburgh PA, USA, (11)U of Arizona, Tucson, AZ, USA, (12)Observatoire de Paris-Paris, France, (13)Istituto di Astrofisica Spaziale, Rome, Italy, (14)Universite de Paris Sud-Orsay, France, (15) University of Washington, (16)Osservatorio Astronomico di Capodimonte, Italy, (17) Cornell University, Ithaca NY

Robert.m.nelson@jpl.nasa.gov, Mail Stop, 183-501, Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena CA 91109, USA

During the Cassini orbital tour Cassini VIMS has observed surface of Titan through spectral 'windows' in its atmosphere where methane, the principal absorbing gas, is transmitting. We have found that the reflectance of a region on Titan's surface (latitude 26S, longitude 78W) increased twofold between July 2004 and March-April 2005. It then returned to the July 2004 level by November 2005. By late December 2005 the reflectance had surged upward again to a new maximum. It then declined for the next three months. Detailed analyses indicate that the brightenings are a surface phenomenon, making these the first changes seen on Titan's surface. The spectral differences between the region and its surroundings rule out the ices of H<sub>2</sub>O, CO<sub>2</sub>, and CH<sub>4</sub> as possible causes. Remarkably, the change is spectrally consistent with the deposition and removal of ammoniated materials. NH<sub>3</sub> has been proposed as a con-

stituent of Titan's interior but not its surface or atmosphere. This transitory  $\text{NH}_3$  spectral signature is consistent with occasional effusion events in which juvenile ammonia is brought to the surface. Its decomposition may feed nitrogen to the atmosphere. The size of the region suggests it may exceed the size of the largest active volcanic areas in the solar system.

This work done at JPL/CALTECH under contract with NASA