



S-wave and density structure along two wide-angle lines in SW-Ireland

F. Hauser(1), B.M. O'Reilly(1), P.W. Readman(1), V.C. Do(1,2) and H.-M. Rumpel(3)

(1) Dublin Institute for Advanced Studies, Dublin 2, Ireland (e-mail:fh@cp.dias.ie), (2) WesternGeco, Buckingham Gate, West Sussex RH6 0NZ, United Kingdom, (3) GGA-Institut, Leibniz Institute for Applied Geosciences, D-30655 Hannover, Germany

VARNET-96 was an international, multidisciplinary project, designed to examine the 'Variscan Front' in southwest Ireland. The seismic experiment consisted of two lines running SSE-NNW. Along line A 8 in-line shots were recorded over a length of ca. 200 km. Line B is ca. 140 km long and recorded 13 in-line shots. Results from ray tracing and travel time inversion of refracted and reflected S-waves indicate a multi-layer crust. It can be sub-divided into a sedimentary layer, an upper crust, a middle crust, a lower crust, and the uppermost mantle. The sedimentary and upper crustal layers show strong vertical and lateral velocity variations along the profiles. Throughout the middle crust velocities increase and reach maximum values of up to 4.1 km/s in the lower crust. The central part of the lower crust along line B shows a strong anomaly. This area was modelled as a low velocity region with values of $V_s=3.90$ km/s. The P- and S-wave crustal velocities define variations in Poisson's ratio, giving better constraints on the likely petrological composition of the crust. This is further constrained by density variations derived from the P- and S- velocities that are correlated with changes in Bouguer gravity across southwest Ireland.