



Mass changes in earth's global water reservoirs

J. Famiglietti (1), D. Chambers (2), F. Frappart (1), S. Nerem (3), M. Rodell (4), S. Swenson (5), I. Velicogna (6,7), J. Wahr (6)

(1) Dept. of Earth System Science, University of California, Irvine, USA, (2) Center for Space Research, University of Texas at Austin, USA, (3) Colorado Center for Astrodynamics Research, Dept. of Aerospace Engineering Sciences, University of Colorado, Boulder, USA, (4) Hydrological Sciences Branch, NASA Goddard Space Flight Center, Greenbelt, MD, USA, (5) Climate and Global Dynamics Division, National Center for Atmospheric Research, Boulder, CO, USA, (6) CIRES and Dept. of Physics, University of Colorado, Boulder, USA, (7) Jet Propulsion Laboratory, California Institute of Technology

Water storage changes in Earth's land, ocean, atmosphere and ice reservoirs vary on time scales ranging from instantaneous to geologic. Until recently, the lack of a consistent, single monitoring framework has hampered efforts to track shorter-term (e.g. monthly, seasonal-interannual (S-I) and decadal-scale) of these storage changes, which are essential for enhanced weather and climate prediction, and for monitoring global change impacts on the hydrologic cycle. The recent launch of the Gravity Recovery and Climate Experiment (GRACE) satellite mission offers a tremendous new opportunity to monitor water storage changes in Earth's major reservoirs. Here we use primarily GRACE observations of time-variable gravity to provide a global assessment of monthly water storage changes in Earth's ocean, ice, land and atmosphere reservoirs. We discuss the emergence of short-term trends for the continents and some major river basins, as well as issues of global water balance closure.