



## **New constraints on hum excitation mechanisms imposed by the discovery of the torsional hum**

**R. Widmer-Schmidrig** (1,2) and **D. Kurrle** (2)

(1) Black Forest Observatory, Schiltach, Germany, (2) Institute of Geophysics, Stuttgart University, Germany.

The recently discovered torsional hum (Kurrle and Widmer-Schmidrig, submitted to GRL, 2008 and same authors EGU 2008) consists of fundamental toroidal modes in the band 3-7 mHz, is persistently present in times devoid of large earthquakes and has an acceleration amplitude of approximately 0.3 nGal which is comparable to the spheroidal hum.

Since the torsional hum signal is dominated by fundamental modes, its excitation should preferentially occur near the Earth's surface, where this mode branch has the largest amplitude. This same argument was already put forth for the source of the spheroidal hum by Kobayashi and Nishida (Science, 1998).

The most intriguing aspect of the torsional hum is that its amplitude more or less equals the spheroidal hum.

The currently favored models for the excitation of the spheroidal hum all involve normal stresses exerted on the Earth's surface either by turbulence in the atmosphere or infragravity waves in the oceans. We can see three mechanisms by which such pressure fields could lead to an excitation of toroidal modes: (1) by the horizontal component of normal stresses on a surface with topography (e.g. mountain torque), (2) by spheroidal-toroidal mode coupling through heterogeneous Earth structure or (3) Coriolis coupling. In all three cases the torsional hum should be at least an order of magnitude smaller than the spheroidal hum. Since this is not the case, new physical mechanisms are called for.

A new mechanism that might go a long way to explain the torsional hum amplitude involves shear stresses exerted on the Earth's surface either by winds over the continents or by currents along the sea floor. If in the mHz-band these vector fields should turn out to have much larger toroidal (divergence free) than poloidal (curl free) horizontal components, they could preferentially excite toroidal over spheroidal modes and selectively excite the torsional hum.

Alternatively, we recall case studies which show that the marine microseisms can consist of Love and Rayleigh waves of equal amplitudes (Rind and Donn, JGR, 1979). Simply by analogy one might speculate that infragravity waves on the shelf also excite the spheroidal and toroidal hum to similar amplitudes.