



Melt flow in a conduit and seismic signals time evolution: A laboratory study

S. Vinciguerra (1), L. Caricchi (2), L. Burlini (3), P. Scarlato (1)

1) Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Roma1, Italy;

2) Institute for Mineralogy and Petrology, ETH, Zürich, Switzerland,

3) Rock deformation Lab, Geological Institute, ETH, Zürich, Switzerland

A variety of seismic signals have been related to fracturing and magma transport in the volcanic edifice. Previous AE studies have provided a first experimental support to this association by reproducing fluid generation and migration while recording seismic signals in a layered sample comprising an olivine-MORB-olivine sandwich.

Here, we developed a new experimental set up, consisting of a melt reservoir beneath a mini conduit, which allowed controlling the physical mechanisms taking place and the related seismic signals during the flow of the melt from the source to the reservoir. The reservoir simulates a natural magma chamber and the mini conduit a volcanic conduit.

Experiments were carried out using an internally-heated Paterson gas apparatus especially designed for the measurements of physical properties of rocks up to 1373K and 300MPa confining pressure. Acoustic emissions were measured during heating of cold pressed synthetic aggregate of MORB powder till complete melting.

High frequency events were recorded at 750-960K, corresponding to glass transitions in the MORB glass. A long lasting low frequency event took place at 1200K, corresponding to melt migration in the conduit, followed by high frequency events, related to brittle intrusive mechanisms due to the magma intrusion in the brittle medium at the top of the conduit. Experiments were also carried out on higher (~2 order of magnitudes) viscosities aggregates of Fish canyon tuffs, and varying the length of the conduit (from 0.5 to 3mm). Results revealed the presence of a new kind of long lasting

signal, made of a cascade of hybrid events with an initial high frequency component, followed by a low frequency signal, which characterizes the intrusion of the silicic magma in the porous reservoir. A direct relationship between seismic waveforms + spectrograms and physical phenomena can be assessed, by scaling length and frequency. These evidences provide solid and well constrained new experimental insight into the association of magma migration in a conduit and related seismic signals. .