



Reconstruction of interdecadal variability of air-sea interaction in the Atlantic 1880-2004: links to atmospheric circulation patterns

S.K.Gulev (1)

(1) P.P.Shirshov Institute of Oceanology, RAS, Moscow

Using 125 years (1880-2004) of Voluntary Observing Ship (VOS) observations from ICOADS we reconstruct surface ocean-atmosphere heat fluxes over the North Atlantic with monthly resolution in time and variable (2-degree to 5-degree) resolution in space. Methodology of reconstruction is based on the homogenization of sampling density, application of the double-exponential distributions of turbulent fluxes for minimizing sampling errors and the use of specially adopted for incomplete data coverage bulk-algorithms. In particular, a multi-regressive approach is used to reconstruct atmospheric humidity, playing important role in estimation surface fresh water fluxes. The methodology was first validated using the time series from VOS and reanalyses for the well sampled last several decades. Further analysis included computation of monthly anomalies of surface fluxes as well as estimation of the subpolar gyre heat and freshwater budgets. These were computed using two-dimensional distributions of surface fluxes in the coordinates of sea-air temperature difference and wind speed. Reconstructed fluxes reveal long-term trends, implying, for example, about 4 W/m² per decade growing sensible heat fluxes in the Labrador Sea and about 2 W/m² per decade secular increase in the Central subpolar gyre. Non-secular signals are represented by the decadal-scale and multidecadal (about 40-50 years variability). Decadal scale signal has a clear association with the NAO-like atmospheric circulation variability during 1880-1915 and after 1955, but has a little association with NAO between 1915 and 1955. The approach formulated allows also for the derivation of the heat energy budgets in different Atlantic regions. These budgets can be alternatively quantified from the oceanographic full-depth sections. Time series of the budget estimates

were derived for 2 large regions (subpolar, mid latitudes) and their association with ocean dynamics and atmospheric circulation anomalies was discussed.