



## **An examination of potential controls on shell Mn/Ca ratios in the calcite of the bivalve *Mytilus edulis***

**P. Freitas** (1, 2), L. Clarke (1), H. Kennedy (1) and C. Richardson (1)

(1) School of Ocean Sciences, Bangor University, UK, (2) Now at the Dept. de Geologia Marinha, INETI, Portugal (pedro.freitas@ineti.pt)

The Mn/Ca ratios in the calcite of marine bivalves have been suggested to reflect both the dissolved and/or particulate Mn concentrations of seawater. However, a clear understanding of what controls shell Mn/Ca ratios is still lacking and a clear quantitative relationship between dissolved and/or particulate Mn and shell Mn/Ca ratios in either calcitic or aragonitic molluscs must be established and validated before any application of a bivalve Mn/Ca palaeoproxy. To study the influence of seawater dissolved and particulate Mn concentrations on bivalve shell calcite Mn/Ca ratios, *Mytilus edulis* specimens were grown in a field experiment in the Menai Strait, U.K., for a one-year period. A single maximum ( $0.54 \mu\text{M}$ ) during spring dominated the annual variation of seawater dissolved Mn concentrations, while seawater particulate Mn concentration was highest (up to  $0.18 \mu\text{M}$ ) during autumn and winter, although smaller increases in particulate Mn during the phytoplankton spring bloom were also observed. In *M. edulis*, shell Mn/Ca ratios of newly precipitated calcite showed a double-peak annual variation with maximum values (up to  $0.19 \text{ mmol/mol}$ ) during early spring and early summer. None of the two maximum of shell Mn/Ca ratios can be explained by an increase in either seawater dissolved or particulate Mn concentrations. Shell Mn/Ca ratios thus were not controlled by dissolved or particulate Mn concentrations. In *M. edulis*, the double-peak seasonal variation of shell Mn/Ca ratios was remarkably similar to the seasonal variation of shell growth rates. However, the influence of shell growth rate on shell Mn/Ca ratios is the opposite of the inverse relationship observed unequivocally between precipitation rate and Mn partition coefficient in synthetic inorganic calcite, and thus must reflect a physiological influence on shell Mn content.