Geophysical Research Abstracts, Vol. 10, EGU2008-A-12012, 2008 SRef-ID: 1607-7962/gra/EGU2008-A-12012 EGU General Assembly 2008 © Author(s) 2008



Detection and mapping of the underground cavities by Geophysical methods: ground penetrating radar seismic waves and electrical imagery. The case of Béni-Mellal city (Morocco)

M. Filahi (1), P. Andrieux (2), A. Tabbagh (2), F. Rejiba (2), A. Najine (1), M. Jaffal (3), A. Casas (4), D. Angelova (5)

(1) Faculty of Sciences and Techniques Beni-Mellal - 23000 Morocco (filahi4@voila.fr), (2) UMR 7619 SISYPHE University Pierre and Marie curie Paris 6 France
(Ep.andrieux@infonie.fr), (3) Faculty of Sciences and Techniques Gueliz Marrakech Morocco (Jaffal_m@yahoo.fr), (4) University of Barcelona faculty of geology department of geochemist 08028 Barcalona (albertc@geo.ub.es), (5) (5) Geological Institute, Bulgarian Academy of sciences 1113 Sofia Bulgaria (doraangelova@hotmail.com)

During the recent years, there has been a growing interest in recording and investigating the network of underground cavities and karsts in many countries around the world. The karstic grounds in Morocco dominate considerable areas. The city of Béni-Mellal has been built on a ground affected by an important network of underground cavities which actually causes numerous buildings collapse. In order to test the efficiency of geophysical methods to localize and map these cavities, the ground penetrating radar campaign with a 200 , 225, 400 MHz antenna, seismic mapping, and 2D electrical tomography measurements have been carried out in the area of the old city (Medina). Prospections at several depths, in the form of sections of resistivity and isochrones maps, allow to identify the resistant zones and the seismic echoes resulting from the interfaces. The analysis of the results showed that these anomalous zones correspond to a broad underground cavity, almost occupying all the width of the area. The methodological tests developed in this study constitute efficient tools to the managers to intervene upstream and to found the preventive means against any risk of collapse. The results clearly reveal the existence of subsurface cavities, the largest one spreads on about fifty square metres.